

NEUTRON STARS WITH A STABLE, LIGHT SUPERSYMMETRIC BARYON

SHMUEL BALBERG¹, GLENNYS R. FARRAR² AND TSVI PIRAN³
NYU-TH-00/09/05

ABSTRACT

If a light gluino exists, the lightest gluino-containing baryon, the S^0 , is a possible candidate for self-interacting dark matter. In this scenario, the simplest explanation for the observed ratio $\Omega_{dm}/\Omega_b \approx 6 - 10$ is that $m_{S^0} \sim 900 \text{ MeV}c^{-2}$; this is not at present excluded by particle physics. Such an S^0 could be present in neutron stars, with hyperon formation serving as an intermediate stage. We calculate equilibrium compositions and equation of state for high density matter with the S^0 , and find that for a wide range of parameters the properties of neutron stars with the S^0 are consistent with observations. In particular, the maximum mass of a nonrotating star is $1.7 - 1.8 M_\odot$, and the presence of the S^0 is helpful in reconciling observed cooling rates with hyperon formation.

Subject headings: dark matter—dense matter—elementary particles—equation of state—stars: neutron

1. INTRODUCTION

The interiors of neutron stars offer a unique meeting point between astrophysics on one hand and nuclear and particle physics on the other. The macroscopic properties of neutron stars, such as mass, radius, spin rate and thermal history depend on the microscopic nature of matter at very high densities. Neutron stars are, in this sense, cosmological laboratories of physics at densities exceeding the nuclear saturation density ($\rho_0 \simeq 0.16 \text{ fm}^{-3}$). One aspect of matter at supernuclear densities is the possible appearance of hadronic degrees of freedom beyond nucleons. The principle candidates are hyperons, meson condensates and a deconfined quark phase (see, e.g., Glendenning 1996, Prakash et al. 1997 for reviews).

In this work we wish to illuminate another possibility, the appearance of an R -baryon. We are motivated by the observation of Spergel & Steinhardt (2000) that the observed properties of dark matter (DM) on scales of order 10-100 kpc are better described by a theory with self-interacting DM than by conventional, weakly interacting DM. Since the required cross-section to mass ratio, $\sigma_{XX}/m_X \sim 1\text{b}/\text{GeV}$, is characteristic of hadronic physics at low energies, a light stable neutral hadron would be a welcome feature of physics-beyond-the standard model, and such particles arise naturally in theories with a very light gluino. In some sense, this is the minimal extension of the standard model needed to produce the required DM particle, and therefore merits examination.

A supersymmetric version of the standard model contains a massless gluino ($\tilde{g}$), the spin-1/2 superpartner of the gluon; therefore it is endowed with a corresponding chiral symmetry called R -invariance. When supersymmetry is broken, the R -invariance may or may not break. If it does not, the gluino mass arises only from loops sensitive to the Higgs vevs (which do break R -invariance) and is naturally of order $100 \text{ MeV}c^{-2}$. Although considerable effort has been devoted to excluding it, a very light gluino remains a possibility – mainly because of the difficulty of making firm, first-principles predictions for non-perturbative phenomena in QCD. The question deserves very serious scrutiny because the existence of a light gluino would resolve several puzzles. These issues and the constraints imposed by experiment are reviewed by Farrar (1998, 2000). See

also Farrar & Gabadadze (2000) and references therein.

The lightest supersymmetric “ R -hadrons” are the spin-1/2 $\tilde{g}g$ bound state (R^0 or glueballino) and the spin-0 baryon $uds\tilde{g}$ bound state, the S^0 . The S^0 is thus a boson with baryon number unity and a strangeness charge -1 . The strong color hyperfine interaction unique to the $uds\tilde{g}$ in a flavor singlet state makes the mass of the S^0 much less than that of all other R -baryons (Farrar 1984; Buccella & Farrar 1985), and probably less than $2 \text{ GeV}c^{-2}$, which would make it stable through conservation of R -parity (Farrar 1996).

In the very early universe the S^0 remains in equilibrium with nucleons via processes such as $R^0 N \leftrightarrow S^0 K$, down to $T \approx 35 \text{ MeV}$ (Farrar, Spergel & Steinhardt 2000; see also in Wandelt et al. 2000). Hence for a suitable S^0 mass, slightly less than the proton mass, standard freezeout estimates lead to a density ratio $\Omega_{S^0}/\Omega_B \approx 6 - 10$, as required if the S^0 is to be the dark matter. To posit a new baryon lighter than a nucleon is quite shocking! But it is known that the color hyperfine force is very strong in this channel, and there are innumerable examples of the inadequacy of phenomenological models of non-perturbative QCD phenomena. The possibility of such a light S^0 should not be dismissed on theoretical grounds, and we must seek observational constraints. One possible astrophysical constraint is whether this new baryon can form in neutron stars, and if so, would it imply unacceptable neutron star properties?

Nucleons cannot convert directly into the S^0 , even if the latter is less massive. Conservation of R -parity requires that R -hadrons are produced in pairs, forbidding direct conversion of individual nucleons into S^0 particles. The weak neutral current process $nn \rightarrow S^0 S^0$ requires simultaneous creation of two units of strangeness ($\Delta S = -2$). Strangeness-changing neutral currents ($\Delta S = \pm 1, \Delta Q = 0$) are highly suppressed in the standard model due to the GIM mechanism. Thus, $nn \rightarrow S^0 S^0$ is doubly GIM suppressed; crude estimates indicate that its rate is of order the inverse Hubble time (Farrar et al. 2000). However processes such as $\Lambda\Lambda \rightarrow S^0 S^0$ ($\Delta S = 0$) and even $\Sigma^- p \rightarrow S^0 S^0$ (charged current, $\Delta S = -1$) are not GIM suppressed, so that if the conditions in interiors of neutron stars allow for hyperons to appear (as is predicted by many works on this issue, see, e.g., Glendenning 1985, Weber & Weigel

¹ Department of Physics, University of Illinois at Urbana–Champaign, 1110 West Green Street, Urbana, IL 61801–3080, sbalberg@astro.physics.uiuc.edu

² Department of Physics, New York University, NY, NY 10003, farrar@physics.nyu.edu

³ Racah Institute of Physics, The Hebrew University, Jerusalem 91904, Israel, tsvi@nikki.phys.huji.ac.il

1989, Knorren, Prakash & Ellis 1995, Schaffner & Mishustin 1996, Balberg & Gal 1997, Huber et al. 1998, Baldo, Burgio & Schulze 2000), these hyperons could serve as a doorway for S^0 formation, which might affect observable properties of neutron stars.

Our goal in this *letter* is to explore whether a light S^0 can form in neutron stars and assess whether the resulting global properties of these stars are compatible with observations. In § 2 we extend an equation of state for high density matter with hyperons to include the S^0 . The equilibrium composition with the S^0 is found in § 3, followed by a discussion of the corresponding properties of neutron stars in § 4.

2. S^0 INTERACTIONS IN HIGH DENSITY MATTER

In order to include S^0 particles in an equation of state of high density matter, we must specify their interactions with other baryons and among themselves. Here we do so in the framework of an effective equation of state (Balberg & Gal 1997, Balberg, Lichtenstadt & Cook 1999), where the potential felt by a single, isospin singlet particle of species x placed in matter composed of species y at a baryon number density ρ_y is

$$V_x(\rho_y) = a_{xy}\rho_y + c_{xy}\rho_y^\gamma + w\rho_y^\theta. \quad (1)$$

In equation (1) a is negative and represents a long range, attractive component in the interaction, c is positive, corresponding to an intermediate range repulsive component, and w is also positive, representing a universal repulsive short range interaction, presumably arising from the quark structure of the baryons. The powers γ and θ must be greater than unity, so that repulsion dominates at high densities.

For nucleons and hyperons, the values of the coefficients can be constrained by experimental data. Since no such data exists for the S^0 , we resort to phenomenological arguments to estimate the magnitude of these terms. *The attractive* component is generally associated with two-pion or two-kaon exchanges. This interaction is strong because the intermediate state in the s -channel can be the same as the initial state, hence the amplitude has a small energy denominator. However the flavor singlet S^0 cannot exchange π or K without becoming a much more massive flavor octet R -baryon (Buccella & Farrar 1985). Thus the long-range attractive interaction is strongly suppressed for the S^0 . A more empirical way to arrive at the same conclusion is to note that if there were a significant long-range attractive force the S^0 would bind to nuclei (and to itself), and nuclei with exotic mass values have not been observed. We therefore assume that the attractive component is negligible for the S^0 -related interactions. *The intermediate repulsive* component may be associated with the exchange of vector mesons, such as the ρ and η . Only the flavor-singlet component of these vector-mesons can couple to the S^0 , so we roughly assume that the magnitude of this term (the value of c in Eq. [1]) is half of its value for the Λ . *The universal short-range* interaction for the S^0 is taken as identical to that of the baryons. Here we use EoS 1 of Balberg et al. (1999), where $\gamma = \frac{4}{3}$ and $\theta = \frac{5}{3}$, so that for the S^0 $a = 0$ and $c = 329.5 \text{ MeVfm}^4$ for all its interactions, and $w = 223.6 \text{ MeVfm}^5$. Below we refer to this choice of parameters along with an S^0 mass of $m_{S^0} = 900 \text{ MeVc}^{-2}$ as our nominal set. Most of our conclusions prove insensitive to these choices, as will be seen below.

⁴ In reality, some additional nucleons would convert to hyperons through quantum and thermal ($T \sim 0.01 \text{ MeV}$ in neutron star cores) fluctuations, and these hyperons would decay to S^0 's. The actual equilibrium composition would be one where the chemical potential of the nucleons is slightly below that of zero-fraction hyperons at every density (in other words, fluctuations would deplete a thin outer layer in phase space of nucleons). Assuming that the probability of fluctuations decreases exponentially with the energy gap, the quantitative effect on the results should be small.

3. COMPOSITION OF NEUTRON STAR MATTER WITH THE S^0

The composition of neutron star matter is determined by chemical equilibrium for all weak-interaction processes. Specifically, these are $B_1 \rightarrow B_2 \ell \nu_\ell$, where B_1 and B_2 are baryons, ℓ is a lepton and ν_ℓ is its corresponding neutrino (or antineutrino). Neutron star matter is well approximated as cold ($T = 0$) and neutrino free ($\mu_\nu = 0$) so that all equilibrium conditions are summarized concisely in the form:

$$\mu_{B_i} = \mu_n - q_{B_i} \mu_e, \quad (2)$$

where μ_{B_i} is the chemical potential of the baryon species B_i , μ_n and μ_e are the neutron and electron chemical potentials, respectively, and q_{B_i} is the electric charge of the baryon B_i . The equilibrium composition at any baryon number density, ρ_B , is then determined by two additional conditions: $\sum_i x_{B_i} = 1$ and charge neutrality, $\sum_i x_{B_i} q_{B_i} + \sum_i x_{\ell} q_{\ell} = 0$, where x_{B_i} and x_{ℓ} denote the fraction per baryon for each baryon and lepton species, respectively.

Equation (2) does not fully apply to the case of the S^0 , since the process $nn \rightarrow S^0 S^0$ is suppressed, even if energetically favorable. The S^0 must maintain a direct equilibrium only with hyperons, i.e., $2\mu_{S^0} = 2\mu_{Y^0}$, and $2\mu_{S^0} = \mu_{Y^-} + \mu_p$, but $2\mu_{S^0} = 2\mu_n$ is *not* required (where Y^0 and Y^- denote a neutral and a negatively charged hyperon, respectively). The significance of this distinction is that a combination of S^0 and nucleons may coexist even while $\mu_{S^0} < \mu_n$, as long as no hyperons are present. On the other hand, a finite hyperon fraction will indirectly impose that $\mu_{S^0} = \mu_n$, so then equation (2) applies to all species.

We emphasize that even if $\mu_{S^0} \leq \mu_n$, nuclear matter cannot “burn” exothermally into S^0 matter. Unlike the case of “strange-quark-stars” (Witten 1984; Alcock, Farhi & Olinto 1986), where a deconfining front converts an entire neutron star into the (hypothetically) lower energy strange quark matter if a seed of of strange quark matter is somehow formed (Olinto 1987), there exists no equivalent stimulant to initiate conversion of nuclear-matter into S^0 -matter, regardless of density or composition.

In Fig. 1a we show for reference the equilibrium composition of neutron star matter as a function of baryon density, using the “standard” equilibrium composition of matter with nucleons and hyperons but no S^0 (hereafter, NH matter) based on EoS 1 of Balberg et al. (1999), assuming that the equation of state is valid up to densities as large as $10\rho_0$. The results, with hyperon formation beginning at about $2\rho_0$, are typical of matter with hyperons (Glendenning 1985; Weber & Weigel 1989; Knorren et al. 1995; Schaffner & Mishustin 1996; Balberg & Gal 1997; Huber et al. 1998; Baldo et al. 2000).

Figure 1b depicts the equilibrium composition when the S^0 is included according to the assumptions of § 2. Due to the low mass of the S^0 , its chemical potential is lower than that of all hyperons, even when the fractions of the latter are infinitesimally small. As a result, all hyperons that form convert to S^0 particles, and the matter consists only of nucleons and S^0 . The composition is determined by the nucleons being in equilibrium with an exactly zero fraction of hyperons⁴. Even though absent from the matter, the hyperons do play a role in determining the amount of S^0 particles which are created.




FIG. 1.— Equilibrium compositions for high density matter for EoS 1 of ref. [1]: (a) matter with hyperons when the S^0 is ignored, (b) including the S^0 with $m_{S^0} = 900 \text{ MeVc}^{-2}$, and (c) same as (b) but with $m_{S^0} = 975 \text{ MeVc}^{-2}$.

Since the S^0 is never in chemical equilibrium with the nucleons (or the zero fraction hyperons), the equilibrium composition would be identical for any $m_{S^0} < 900 \text{ MeVc}^{-2}$. In fact, for the nominal interactions, the same composition is also found for $m_{S^0} \lesssim m_p$. A qualitative change is found when m_{S^0} is larger: in Fig. 1c we show the equilibrium composition for nominal S^0 interactions, but with $m_{S^0} = 975 \text{ MeVc}^{-2}$. At low and intermediate densities the composition is identical to that of the $m_{S^0} = 900 \text{ MeVc}^{-2}$ case, but beginning at a density of 0.9 fm^{-3} the larger mass of the S^0 makes it favorable for a finite fraction of hyperons to coexist in the matter. The general equilibrium condition (Eq. [2]) is now satisfied for all particle species. Owing to the S^0 having only repulsive interactions, the hyperons achieve chemical equilibrium with it even though it has a lower mass and no kinetic energy. Moreover, for larger densities, it becomes increasingly favorable to reduce the S^0 fraction as the equilibrium composition gradually approaches that of NH matter. With similar calculations we find that for nominal interactions, if $m_{S^0} \simeq 1030 \text{ MeVc}^{-2}$ the S^0 appears only after some hyperons already exist in the matter, and if $m_{S^0} \gtrsim 1060 \text{ MeVc}^{-2}$ S^0 formation is suppressed for the entire range of densities.

4. EQUATION OF STATE AND MASS SEQUENCES

We now turn to examine the equation of state (pressure vs. mass-energy density, hereafter EoS) of high density matter with the S^0 . These are shown in Fig. 2, including the models used for Fig. 1 and also a model labeled N for nuclear matter,




FIG. 2.— Equation of state for matter of the compositions shown in Fig. 1. Matter with the S^0 : nominal values (see text) - thick solid line; nominal interactions and $m_{S^0} = 975 \text{ MeVc}^{-2}$ - dashed line; nominal mass and increased repulsion - dot-dashed line. Also shown are the equations of state for nucleon and hyperon matter - N+H, and nuclear matter - N.

where hyperon (and therefore also S^0) formation is ignored.

The hierarchy of EoSs in Fig. 2 is mostly determined by the number of available species in each model. Nuclear matter naturally has the stiffest EoS, followed by matter with the S^0 . Even though the S^0 , being a boson, does not exert any Fermi pressure, the fact that four species of hyperons are suppressed due to its appearance makes such matter stiffer than NH matter, which is clearly the softest. An exception occurs just above the threshold density for hyperon formation, where NH matter includes only a single hyperon species (the Σ^-), so that the contribution of its Fermi pressure does make NH matter stiffer.

Although the mass of the S^0 does not affect its contribution to the pressure, it does contribute to the mass energy-density. Hence a lower m_{S^0} yields a stiffer EoS, while a larger value yields a softer one. For $m_{S^0} = 975 \text{ MeVc}^{-2}$ such softening is enhanced at a density of $\sim 1.8 \times 10^{15} \text{ gcm}^{-3}$, when hyperons begin to accumulate; as the S^0 fraction is depleted this EoS gradually approaches that of NH matter. It is noteworthy that increasing the repulsive power of the S^0 does not lead to significant stiffening of the EoS. We demonstrate this with another model shown in Fig. 2 when $m_{S^0} = 900 \text{ MeVc}^{-2}$ but $c = 494.5 \text{ MeV fm}^4$ (50% larger than the nominal value). Magnified repulsion leads to an increase of the S^0 chemical potential, which in turn enhances hyperon formation. The resulting composition of this model is quite similar to that of Fig. 1c, and the EoS, while slightly stiffer than nominal until hyperons appear, is then softened. We emphasize that this is a manifestation of the ability of hyperons to exert “pressure control” (Balberg & Gal 1997; Balberg et al. 1999), since the extent of their formation tracks the dependence of particle interactions on density.

Given an EoS, the structure of a spherical, non-rotating neutron star is found by numerically integrating the Tolman-Oppenheimer-Volkoff equations of general relativistic hydrostatic equilibrium (see, e.g., Misner, Thorne & Wheeler 1973). The resulting mass sequences (stellar mass as a function of the assumed central density) for the EoSs of Fig. 2 are given in Fig. 3. All sequences are identical up to about $1.1 M_\odot$, where the mass sequences diverge according to the hierarchy of EoSs. The maximum mass found for matter with the S^0 is $1.81 M_\odot$ for the nominal case, $1.68 M_\odot$ for $m_{S^0} = 975 \text{ MeVc}^{-2}$,




FIG. 3.— Nonrotating neutron star sequences for the five equations of state shown in Fig. 2.

and $1.78 M_{\odot}$ for $m_{S^0} = 900 \text{ MeV}c^{-2}$ but enhanced S^0 repulsive interactions. In both latter cases, hyperons coexist with the S^0 only in the largest mass stars, those of mass closer than $0.1 M_{\odot}$ to the maximum mass in each case. We note that for EoS 1, the maximum masses for nuclear matter and NH matter are ~ 2.18 and $\sim 1.57 M_{\odot}$ respectively.

At present, the primary observational constraint on high density equations of state is that the maximum mass must be at least $1.44 M_{\odot}$, the inferred mass of PSR 1913+16. All EoSs discussed above, with and without the S^0 , are acceptable. This limit does allow to rule out very soft EoSs, such as would arise if the S^0 had a low mass *and* an attractive component in its interactions. Such interactions would make a pair of S^0 similar to the hypothetical H dibaryon - a six quark flavor singlet, boson (Jaffe 1977a, 1977b) - whose interactions are supposed to be similar to those of two Λ hyperons combined. The EoS for matter with the H would indeed be too soft to sustain a $1.44 M_{\odot}$ star, unless its mass is significantly above $2m_{\Lambda}$ (Glendenning & Schaffner-Bielich 1998). The expected strongly repulsive interactions of the S^0 are thus crucial for allowing its existence in neutron stars.

5. CONCLUSIONS AND DISCUSSION

Our primary conclusion is that the existence of a light ($m_{S^0} \lesssim m_p$) S^0 particle is consistent with the fundamen-

tal observed properties of neutron stars, namely, their inferred masses. This conclusion applies for a wide range of models for high density matter with the S^0 . In particular, the maximum mass of a static neutron star under nominal assumptions regarding the S^0 is found to be about $1.8 M_{\odot}$. We note that since it suppresses hyperon accumulation, the appearance of the S^0 yields a stiffer equation of state and higher maximum mass than obtained with nucleons and hyperons without S^0 . Therefore, identification of a relatively high mass neutron star would be easier to reconcile with an S^0 component than in its absence (where hyperons accumulate undisturbed). Some indication for the existence of such massive neutron stars is implied by recent analyses, although uncertainties are still significant (van Kerkwijk 2000). We emphasize that *current* data on neutron star masses is not restrictive enough to decisively constrain the properties of high density matter.

The presence of S^0 particles could have a significant and welcome effect on the cooling rate of neutron stars, if they do indeed replace hyperons in the cores. Hyperons can participate in the direct Urca cooling process ($B_1 \rightarrow B_2 e \nu$) while the S^0 cannot. The threshold fraction required for hyperons to drive direct Urca cooling processes is of the order of 0.01 (Prakash et al. 1992), so neutron stars with hyperons are susceptible to rapid cooling, which seems inconsistent with observations (Schaab et al. 1996). Hyperon direct Urca cooling can be partially suppressed if the Λ hyperons pair to a $^1S_0$ superfluid state (Schaab, Balberg & Schaffner-Bielich 1998), but the extent of suppression is sensitive to the details of the equilibrium composition and gap energy (Page et al. 2000). On the other hand, S^0 formation would prevent the hyperon direct Urca, or restrict it only to the highest mass stars, so that very moderate cooling rates are attainable. Future measurements of surface thermal emission from young neutron stars with the *CHANDRA* and *XMM* satellites may lead to stronger constraints on cooling rates; if so this will yield valuable insight into the internal structure of high density matter.

The research of GRF is supported in part by NSF-PHY-99-96173. The idea that the S^0 could be a dark matter candidate, and constraints on such a scenario, were developed by GRF in collaboration with D. Spergel and P. Steinhardt; that work will be reported in greater depth elsewhere (Farrar et al. 2000); a brief description can be found in (Wandelt et al. 2000). SB and TP thank NYU for hospitality which facilitated this research.

REFERENCES

- Alcock, C., Farhi, E., & Olinto, A. V. 1986 *ApJ*, 310, 261
 Balberg, S., & Gal, A. 1997, *Nucl. Phys. A*, 625, 435
 Balberg, S., Lichtenstadt, I., & Cook, G. B. 1999, *ApJS*, 121, 515
 Baldo, M., Burgio, G. F., & Schulze, H. J. 2000 *Phys. Rev. C*, 61, 055801
 Farrar, G. R. 1984 *Phys. Rev. Lett.*, 53, 1029
 Buccella, F., Farrar, G. R., & Pugliese, A. 1985, *Phys. Lett. B* 153, 311
 Farrar, G. R. 1996, *Phys. Rev. Lett.*, 76, 4111
 Farrar, G. R. 1998, *Nucl. Phys. Proc. Suppl.* 62, 485
 Farrar, G. R. 2000, (NYU-TH-00/09/08) in preparation
 Farrar, G. R., & Gabadadze, G. 2000, (NYU-TH-00/09/06) in preparation
 Farrar, G. R., Spergel, D. N., & Steinhardt, P. J. 2000 in preparation
 Glendenning, N. K. 1985, *ApJ*, 293, 470
 Glendenning, N. K. 1996, *Compact Stars*, (New York: Springer)
 Glendenning, N. K., & Schaffner-Bielich, J. 1998 *Phys. Rev. C*, 58, 1298
 Huber, H., Weber, F., Weigel, M. K., & Schaab, Ch. 1998, *Int. J. Mod. Phys. E* 7, 301
 Jaffe, R. L. 1977, *Phys. Rev. Lett.*, 38, 195
 — 1977b, *Phys. Rev. Lett.*, 38, 617
 Knorren, R., Prakash, M., & Ellis, P. J. 1995, *Phys. Rev. C*, 52, 3470
 Misner, C. W., Thorne, K. S., & Wheeler, J. A. 1973, *Gravitation*, Freeman, San Francisco.
 Olinto, A. V. 1987, *Phys. Lett. B* 192, 71
 Page, D., Prakash, M., Lattimer, J. M., & Steiner, A. 2000, *Phys. Rev. Lett.*, 85, 2048
 Prakash, M., Prakash, M., Lattimer, M. J., & Pethick, C. J. 1992, *ApJ* 390, L77
 Prakash, M. 1994, *Phys. Rep.*, 242, 297
 Prakash, M. et al. 1997, *Phys. Rep.*, 280, 1
 Schaab, Ch., Weber, F., Weigel, M. K., & Glendenning, N. K. 1996, *Nucl. Phys. A*, 605, 531
 Schaab, Ch., Balberg, S., & Schaffner-Bielich, J. 1998, *ApJ*, 504, L99
 Schaffner, J., & Mishustin, I. N. 1996, *Phys. Rev. C*, 53, 1416
 Spergel, D. N., & Steinhardt, P. J. 2000, *Phys. Rev. Lett.*, 84, 3760
 van Kerkwijk, M. H. 2000, Neutron Star Mass Determinations, *To appear in "Proc. ESO Workshop on Black Holes in Binaries and Galactic Nuclei"*, (L. Kaper, E. P. J. van den Heuvel, and P. A. Woudt, Eds.). Springer-Verlag, Berlin (2000) (astro-ph/0001077)
 B. Wandelt et al. 2000, astro-ph/0006344.
 Weber, F., & Weigel, M. K. 1989, *Nucl. Phys. A*, 505, 779
 Witten, E. 1984, *Phys. Rev. D*, 30, 272